An impressionistic landscape painting featuring a row of trees in the foreground, rolling green hills in the middle ground, and a sky with soft, blended colors of pink, orange, and blue. The style is characterized by visible brushstrokes and a focus on light and color over fine detail.

YEARS OF LIFE LOST DUE TO COVID-19 IN SERBIA: REGISTERED DISTRICT-LEVEL DIFFERENCES

Institute of Social medicine
Faculty of Medicine
University of Belgrade

Presenter:
Teaching Assistant
Aleksandar Stevanović, MD

AUTHORS

Aleksandar Stevanović^{1,2}

Milena Šantrić-Milićević^{1,2}

Jovana Todorović^{1,2}

Stefan Mandić-Rajčević^{1,2}

Nataša Rosić³

Gordana Bjelobrk⁴

Elena von der Lippe

Brecht Devleesschauwer⁶

¹**Institute of Social Medicine**, Faculty of Medicine, University of Belgrade, Serbia

²**Center – School of Public Health**, Faculty of Medicine, University of Belgrade, Serbia

³**City Institute of Public Health Belgrade**, Belgrade, Serbia

⁴**Statistical Office of the Republic of Serbia**, Belgrade, Serbia

⁵**Department of Epidemiology and Health Monitoring**, RKI, Berlin, Germany

⁶**Department of Epidemiology and Public Health**, Sciensano, Brussels, Belgium

CONTENTS

1. Background
2. Methods
3. Results
4. Conclusion
5. Discussion?



BACKGROUND

COVID-19 IN SERBIA

Official onset:
March 6, 2020

Up to December 31, 2022:

- 2,446,253 infection cases
- 17,519 death cases

(data published by IOPH Serbia)

Vital statistics are provided by the Statistical Office of the Republic of Serbia (SORS). This includes the mortality data disaggregated by cause of death.

METHODS

METHODS: DATA

Disaggregated, anonymized information on individual **COVID-19 death cases:**

- **U07.1** – SARS-CoV-2 infection laboratory-confirmed
- **U07.2** – SARS-CoV-2 infection clinically suspected

Sociodemographic characteristics of each case:

- Age
- Sex
- Time and place of death
- Residential Area
- Marital Status
- Educational Attainment
- Activity

METHODS: YLL

- **Years of Life Lost (YLL)** were calculated for each death case individually by assigning the life expectancy of a corresponding gender-age group
- For **the remaining life expectancy**, we used national values provided by the Statistical Office of the Republic of Serbia
- All values were summed up to get the **cumulative YLL number** for each populational group of interest

METHODS: ANALYSIS

- COVID-19 YLL was described **as absolute, average, and relative numbers** (percent and rates per 100,000 population)
- **Pearson's χ^2 test** was used to explore differences among different population groups (men vs. women, urban vs. non-urban settlements, and district-level)

Управни окрузи



30 districts in total (NUTS3)

Data from **25 districts** provided by the Statistical Office of the Republic of Serbia

METHODS: ANALYSIS




1:1,840,000


Урађено у Републичком заводу за статистику
Извор: Графика подлога Републичког геодетског завода

RESULTS


Average number of YLLs per COVID-19 death case in districts of Serbia (2020-2021)




COVID-19 YLLs rate for 2020 & 2021 (per 100,000 population)



Place of COVID-19 death cases (%)



Confirmation of COVID-19 death cases (U07.1 vs U07.2)



CONCLUSION

CONCLUSION

- A notable variance in premature mortality due to COVID-19 in Serbia has been registered at the district level
- Along with the average YLL number, there are significant disparities in the share of deaths that occur outside of hospitals or have no laboratory confirmation
- Additional analysis should try to explain these differences

THANK YOU! 😊 QUESTIONS?